

## PREDICTION OF BLEEDING IN PATIENTS UNDERGOING PERCUTANEOUS CORONARY INTERVENTION: A RISK SCORE USING PRE-PROCEDURAL CLINICAL VARIABLES

i2 Poster Contributions

Ernest N. Morial Convention Center, Hall F

Tuesday, April 05, 2011, 9:30 a.m.-10:45 a.m.

Session Title: Vascular Access, Closure Devices, Complications

Abstract Category: 24. Vascular Access, Closure Devices and Complications

Session-Poster Board Number: 2518-531

Authors: *Megan Coylewright, Paul Sorajja, Ryan J. Lennon, Malcolm R. Bell, Henry H. Ting, Charanjit S. Rihal, John F. Bresnahan, Mandeep Singh, David R. Holmes, Jr., Mayo Clinic, Rochester, MN*

**Background:** Antithrombotic therapies have markedly reduced ischemic complications in patients undergoing percutaneous coronary intervention (PCI). However, their use increases risk of peri-procedural bleeding, which is associated with higher morbidity and mortality. We investigated the use of pre-procedural risk factors to identify patients at heightened risk of bleeding following PCI.

**Methods:** Consecutive patients who underwent PCI via femoral approach at Mayo Clinic from 2000-2009 were included. Major bleeding events included intracranial, retroperitoneal, or gastrointestinal bleeding; pseudoaneurysm; or any bleeding which required transfusion, surgery or prolonged hospital stay. A risk score was created based on coefficients proportional to the parameter estimates from the logistic regression model. It was internally validated via bootstrapping to correct for optimism of the model.

**Results:** Overall, 949 (7.7%) of 12,256 patients had a major bleed. The predicted risk of major bleeding ranged from 1% to >50%. (see Figure 1) Age, gender, recent myocardial infarction, shock, creatinine, body mass index, sheath size and use of glycoprotein IIb/IIIa inhibitors were independent predictors of bleeding. The optimism corrected c-index of the model was 0.74; the Hosmer-Lemeshow goodness-of-fit test was non-significant ( $p=0.09$ ).

**Conclusion:** Pre-procedural patient characteristics may be used to identify patients at high risk of bleeding following PCI and assist in efforts to reduce risk.

